

# Highly processed foods can be considered addictive substances based on established scientific criteria

Ashley N. Gearhardt<sup>1</sup>  | Alexandra G. DiFeliceantonio<sup>2,3</sup>

<sup>1</sup>Department of Psychology, University of Michigan, Ann Arbor, MI, USA

<sup>2</sup>Fralin Biomedical Research Institute at VTC, Roanoke, VA, USA

<sup>3</sup>Department of Human Nutrition Foods, and Exercise, Virginia Polytechnic Institute and State University, Blacksburg, VA, USA

## Correspondence

Ashley N. Gearhardt PhD, Department of Psychology, University of Michigan, Ann Arbor, MI, USA.

Email: [agearhar@umich.edu](mailto:agearhar@umich.edu)

## Funding information

A.G. DiFeliceantonio is an iTHRIV Scholar. The iTHRIV Scholars Program is supported in part by the National Center for Advancing Translational Sciences of the National Institutes of Health, Grant/Award Numbers: UL1TR003015, KL2TR003016

## Abstract

**Background:** There is growing evidence that an addictive-eating phenotype may exist. There is significant debate regarding whether highly processed foods (HPFs; foods with refined carbohydrates and/or added fats) are addictive. The lack of scientifically grounded criteria to evaluate the addictive nature of HPFs has hindered the resolution of this debate.

**Analysis:** The most recent scientific debate regarding a substance's addictive potential centered around tobacco. In 1988, the Surgeon General issued a report identifying tobacco products as addictive based on three primary scientific criteria: their ability to (1) cause highly controlled or compulsive use, (2) cause psychoactive (i.e. mood-altering) effects via their effect on the brain and (3) reinforce behavior. Scientific advances have now identified the ability of tobacco products to (4) trigger strong urges or craving as another important indicator of addictive potential. Here, we propose that these four criteria provide scientifically valid benchmarks that can be used to evaluate the addictiveness of HPFs. Then, we review the evidence regarding whether HPFs meet each criterion. Finally, we consider the implications of labeling HPFs as addictive.

**Conclusion:** Highly processed foods (HPFs) can meet the criteria to be labeled as addictive substances using the standards set for tobacco products. The addictive potential of HPFs may be a key factor contributing to the high public health costs associated with a food environment dominated by cheap, accessible and heavily marketed HPFs.

## KEYWORDS

Addiction criteria, carbohydrates, compulsion, fat, food addiction, highly processed foods, nicotine, reinforcement, smoking, tobacco

## INTRODUCTION

There is evidence that an eating phenotype exists that reflects the hallmarks of addiction (e.g. loss of control over intake, intense cravings, inability to cut down and continued use despite negative consequences) [1]. Based on meta-analyses, approximately 14% of adults and 12% of children exhibit this addictive-like eating phenotype, commonly called food addiction [2, 3]. Although some have questioned

the utility of applying an addiction framework to food intake [4–7], food addiction is associated with mechanisms implicated in other addictive disorders (e.g. impulsivity, reward dysfunction and emotion dysregulation), as well as a lower quality of life and a poorer response to weight-loss treatments [1, 8, 9]. Controversy exists surrounding the role of the food in triggering this addictive-like eating phenotype. Some propose that it is the act of eating regardless of the type of food consumed that is addicting [10], or that while the type of food is

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Addiction* published by John Wiley & Sons Ltd on behalf of Society for the Study of Addiction.

important, it is impossible to classify food as addictive due to the complex nature of foods and the lack of a single addictive agent/compound [4, 5]. Food is necessary for survival and a key evolutionary pressure that has shaped reward and motivation systems across species [11, 12]. Addictive drugs that deliver high doses of reinforcing substances through rapid delivery systems tap into these systems, potentially activate them and can lead to maladaptive patterns of behavior [13]. Highly processed foods (HPFs) are evolutionarily novel products made possible through modern food technology that provide refined and rapidly delivered primary reinforcers, specifically calories in the form of refined carbohydrates and added fats [1, 14–16]. The debate that remains concerns whether a refined and optimized delivery system of calories can produce comparative effects to a refined and optimized delivery system of addictive drugs.

The ability to resolve the debate about whether certain foods are addictive is hindered by a lack of identified scientifically based criteria with which to evaluate the addictiveness of certain foods. In contrast, there is a general consensus around the criteria for identifying whether someone is exhibiting an addictive phenotype [17], which has allowed for clearer criteria to guide the investigation into whether certain individuals exhibit addictive-like eating [2, 3]. There is no comparable standard for evaluating if a substance is addictive, which contributes to the conflicting explanations for why certain foods are (or are not) addictive [18].

To allow for progress on this debate, we propose a set of scientifically based criteria for the evaluation of whether certain foods are addictive. Specifically, we propose the primary criteria used to resolve one of the last major controversies over whether a substance, tobacco products, was addictive. In 1988, the Surgeon General (SG) released a report outlining scientific evidence that tobacco products were addictive. This report acknowledged that no singular criterion is sufficient to determine that a substance is addictive, and frequent (or even excessive) intake alone does not necessarily indicate addiction [19]. Rather, the SG report laid out a set of primary criteria that were identified as being necessary and sufficient to establish the addictive nature of tobacco products: (1) they trigger compulsive use, (2) they have psychoactive effects and (3) they are reinforcing [19]. A recent review confirmed that the main conclusions of this report have held up to over 30 years of scientific evaluation with some important updates based on advances in addiction science, specifically that tobacco products (4) trigger strong urges and craving [20]. The 1988 SG report was a watershed moment that bolstered the scientific recognition of tobacco products as addictive and shifted the public's perception of their addictive nature [20]. This helped to lay the groundwork for the development of new treatments that target addictive mechanisms and effective public health campaigns about the risks of tobacco products.

In this analysis, we consider the evidence that HPFs (i.e. foods with refined carbohydrates and/or added fats) [1, 14–16] can meet all the primary scientific criteria from the 1988 SG report on the addictiveness of tobacco (updated to include the ability to trigger urges and cravings). We propose that the 1988 SG report criteria used for tobacco products are ideal for evaluating the

addictive nature of HPFs, as both these products are legal, easily accessible, inexpensive, lack an intoxication syndrome and are major causes of preventable death. Based on the established scientific criteria on the addictiveness of tobacco, we conclude that HPFs can meet all the criteria for an addictive substance. Below, we outline the evidence for this conclusion, highlight scientific gaps, provide a roadmap for future research and consider the implications of labeling HPFs as addictive.

## PARALLELS IN THE INDUSTRIALIZATION OF FOOD AND CIGARETTE PRODUCTION

A number of terms are used to refer to foods that are prone to excessive intake (e.g. highly processed, hyperpalatable and ultra-processed), and there is a significant debate about which approach is most appropriate [21–24]. We will focus upon food categorized as HPF based on the presence of refined carbohydrates and/or added fats, as these ingredients have been most implicated in addictive-like eating [1, 14–16]. Commonly consumed HPFs include carbonated soft drinks, sweet or savory snacks, chocolate, candies, ice cream, cakes, cookies, bread and pizza (among many others) [25]. Although processed ingredients (e.g. white flour, sugar and butter) have long been available for purchase to allow for the creation of homemade HPFs (e.g. freshly baked bread and homemade cookies), a marked increase in cheap, easily accessible and heavily marketed industrialized HPFs began in the 1980s and preceded increases in obesity and diet-related disease (e.g. Type II diabetes) [26]. The dominance of industrial HPFs is the result of these products being (1) inexpensive due to the use of cheap but potent industrial substances (e.g. high fructose corn syrup and trans fats); (2) highly accessible and convenient due to additives that allow them to stay self-stable and ready-to-eat; (3) enhanced through additives that amplify somatosensory properties related to smell, taste and texture through substances not available to the home cook (e.g. flavorants, monosodium glutamate and guar gum); and (4) the focus of marketing campaigns that create positive associations and brand loyalty from a young age [25–27].

There are clear parallels between industrial HPFs and industrial tobacco products. Although dried tobacco leaves have been consumed through smoking or chewing since at least AD 800, the development of industrial cigarette-rolling machines in the 1880s revolutionized the tobacco industry [28]. Cigarettes could now be rolled at more than 10 000 a minute (compared to just a few a minute when hand-rolled). This technological shift allowed for the mass production of inexpensive industrial cigarettes and the development of the modern tobacco industry [28]. As with industrial HPFs, industrial cigarettes are convenient, accessible and heavily marketed [29]. Although hand-rolled cigarettes made from processed tobacco can still be addictive and damaging to health, it was the marked increase in cheap, convenient, accessible and heavily marketed industrial tobacco cigarettes that led to a > 1000% increase in cigarette consumption and a marked increase in smoking-related disease (e.g. lung cancer) [30]. Similarly, while home-made HPFs that include processed

ingredients (e.g. sugar and fat) can be consumed addictively and damage health, the rise of cheap, accessible, convenient and heavily marketed industrial HPFs in the food supply is the major driver of excessive food intake, obesity and diet-related diseases [25–27]. In contrast, the intake of minimally processed foods (MPFs) that promote health, such as fruit, vegetables and legumes, fails to reach recommended levels despite well-funded public health campaigns to promote their consumption [31].

## PRIMARY SCIENTIFIC CRITERIA FOR EVALUATING THE ADDICTIVENESS OF A SUBSTANCE

### Criterion 1: compulsive use

We will now consider whether HPFs meet all criteria outlined in the SG report as necessary and sufficient to identify tobacco products as addictive. The first criterion in the SG report is the ability of the substance to trigger compulsive use, defined by ‘drug-seeking and drug-taking behavior that is driven by strong, often irresistible urges. It can persist despite a desire to quit or even repeated attempts to quit’ [19]. Intake alone is not sufficient to identify compulsive use, as only a subset (one-third) of tobacco smokers develop addictive patterns of use [32]. Compulsive use for tobacco in the SG report was demonstrated by evidence that most smokers would like to quit, but most were unable to do so. The report notes that compulsive nature is most clear in extreme cases where individuals who are experiencing significant smoking-related disease (e.g. cancer and cardiovascular disease) continue smoking [19].

It is clear that HPFs are capable of triggering compulsive use. Even in the face of significant diet-related health consequences (e.g. diabetes and cardiovascular disease), the majority of patients are unable to adhere to medically recommended dietary plans that require a reduction in HPF intake and a commonly cited obstacle for low dietary adherence is cravings for HPFs [33–37]. Failure in response to gastric bypass surgery (i.e. a significant surgery that results in the division of the stomach to treat severe obesity) provides an extreme case of compulsive HPF intake. Approximately 20–50% of individuals who undergo this surgery regain a significant amount of weight [38] and excessive HPF intake is a factor in this weight regain [38–40]. This intake persists despite HPFs triggering immediate aversive physical symptoms (e.g. cramping, vomiting and diarrhea) when consumed after surgery [38–40].

Binge eating (i.e. the experience of a loss of control when consuming a large amount of food) also provides evidence of compulsive HPF intake [41]. The type of food predominantly consumed in binges are HPFs, whereas MPFs (such as fruits and vegetables) are less likely to be consumed during binges [42–44]. Higher dietary intake of MPFs, including high-fat nuts, is inversely associated with binge eating disorder [45]. In contrast, higher dietary intake of HPFs, such as sweets and baked goods, is positively associated with binge eating disorder [45]. Similarly, rodents will risk aversive experiences

(e.g. electric shock) to consume industrially produced cheesecake, pound cake, frosting and chocolate when other calorie sources (i.e. standard chow) are easily available to them [46]. Rats even show greater resistance to punishment (foot shock) when working for a ‘preferred pellet’ reward (i.e. chow with added corn syrup) than when methamphetamine is used as the reinforcer [47]. In sum, HPFs, but not MPFs, appear to meet the criterion of triggering compulsive intake consistent with addictive substances.

### Criterion 2: psychoactivity

Another primary criterion outlined in the SG report is that the substance needs to cause psychoactive effects [19]. Psychoactivity was defined as ‘produces transient alterations in mood that are primarily mediated by effects in the brain’ [19]. The ability of tobacco to alter mood is more subtle than intoxicating substances, such as opioids and alcohol. However, tobacco products can cause detectable subjective increases in pleasure and reductions in negative affect [19, 48]. These mood-altering effects are related to the ability of tobacco products to deliver high doses of nicotine rapidly to the brain [19]. It is important to note that at the time of the SG report exact knowledge of how nicotine impacted on the brain, particularly in humans, was limited and mainly focused upon the ability of the drug to bind to nicotinic receptors and increase glucose utilization in the rat brain [19]. Those studies identified a host of brain regions, now known to be dense in nicotinic receptors, but of those, only the medial habenula and ventral tegmental area are established as key mediators of nicotine self-administration and use [49]. In the SG report, the ability of centrally acting nicotinic antagonist to alter tobacco smoking also provided evidence about the important role of the brain in the use of tobacco products [19].

Advances in addiction science the last 30 years have yet to lead to the discovery of a specific biomarker to identify whether or not a substance is addictive, but there is a consensus that all addictive substances increase dopamine in the striatum [50]. Relative to dopamine agonists such as amphetamine, which can increase striatal dopamine release by 1000% [51], nicotine administration causes more modest increases in dopamine efflux (150–250% over baseline), which is similar to other addictive drugs such as ethanol (also 150–200% over baseline) [51, 52]. However, despite this lower magnitude, nicotine is still capable of triggering compulsive intake and changing mood. Since the SG report, it has been demonstrated that the amount of subjective pleasure experienced in response to tobacco products is weakly associated with relapse and may be less central to its addictive nature [20]. In contrast, the ability of tobacco products to trigger strong urges and cravings appears to be a bigger mediator of addictive patterns of intake, which will be reviewed in more detail in Criterion 4 [20].

Based on the standards above, HPFs can be considered psychoactive substances. HPFs are capable of increasing positive affect and reducing negative affect [16, 53–57]. For example, intake of white chocolate and 38% cocoa chocolate are associated with ‘euphoria’

ratings on a measure of psychoactive drug effects of 7.0 and 6.4, respectively [53], which is similar to the 'euphoria' scores on the same measure after the administration of 1.5 mg of intravenous nicotine to smokers (score of ~6) and much higher than scores by non-smokers (~1) [48]. Intake of HPFs, relative to MPFs, is associated with greater subjective experiences of enjoyment (21.69 points higher on a visual analog scale) and the magnitude of this effect is greater for individuals with addictive-like eating [57]. Further, HPF intake is often motivated by a desire to alter mood (e.g. experience pleasure and reduce negative affect) rather than to address homeostatic needs and this tendency is associated with addictive eating [1, 58]. Further research is warranted to directly compare the magnitude of the psychoactive effects of a wider range of HPFs to known addictive substances and MPFs, but existent research supports the psychoactive nature of HPFs.

Regarding the brain, HPFs and their components increase dopamine in the striatum at a similar magnitude as nicotine when delivered orally (150–200%) [59–62]. These substances increase striatal dopamine (~150%) and dopaminergic firing rates even when oral somatosensation is bypassed and nutrients are delivered directly to the gut [63–66]. Whereas MPFs are predominantly high in either carbohydrates (e.g. fruit) or fat (e.g. nuts), HPFs are unique in their tendency to rapidly deliver a combination of refined carbohydrates and fat. The combination of refined carbohydrates (such as sugar) and fat appears to have a supra-additive effect on reward encoding in the striatum in humans [67]. Opioid antagonists are also capable of reducing HPF (including chocolate candies and cookies) intake in humans [68–71]. In animal models, opioid antagonists reduce consumption of chocolate chip cookies, but not laboratory chow [68, 72], which further highlights the importance of reward systems in the brain in driving HPF intake. As with tobacco, the experience of subjective liking of foods may be less central to their tendency to maintain compulsive intake, but rather their ability to trigger strong urges and cravings may be more central to their addictive potential [73, 74]. However, there is sufficient evidence to label HPFs as psychoactive substances based on the SG report.

### Criterion 3: reinforcing

The final criterion in the SG report was that tobacco was a reinforcing substance, as defined by 'being sufficiently rewarding to maintain self-administration' [19]. Clearly, humans will self-administer tobacco products, although not all humans find tobacco products reinforcing [48, 75]. Nicotine was identified as a key factor in the reinforcing nature of tobacco products, as animals would self-administer nicotine, work to gain access to nicotine, prefer places where nicotine was administered and conditioned cues paired with nicotine became secondary reinforcers [19]. Compared to other addictive drugs (such as cocaine), nicotine was a relatively weak reinforcer and was only self-administered under a narrow range of conditions (e.g. intermittent reinforcement schedules, food-restriction, combined with food delivery, paired with cues) [19]. However, this level of evidence was sufficient for the SG report to conclude that tobacco products were reinforcing due to their ability to deliver nicotine.

Scientific evidence suggests that the reinforcing nature of HPFs is high. Both adults and children will self-administer HPFs (e.g. potato chips, candy and cookies) even when satiated [76, 77]. The tendency to consume MPFs when satiated appears to be lower. For example, when children are provided with post-meal access to a MPF (fruit) rather than an HPF (sweet/savory snack), energy intake is reduced by an average of 60% [78]. Both adults and children will work on an operant responding paradigm (a classic measure of reinforcement) to gain access to HPF relative to non-food reinforcers [79]. Adults, adolescents and children who find HPFs more reinforcing (as indicated by a greater willingness to work to gain access to them) are more likely to have obesity and are prone to weight gain [80–83]. In contrast, the reinforcing value of MPF (e.g. fruit, vegetables, cottage cheese) is not associated with higher body weight in adults, adolescents and children [80, 84, 85]. Daily exposure to HPFs (e.g. candy, cookies, chips) appears to sensitize the reinforcing value of these foods (as indicated by an increased willingness to work to gain access to HPFs over time) and larger portions of HPFs lead to greater sensitization [84–87]. In contrast, daily exposure to MPFs, such as vegetables and fruits, does not sensitize reinforcement and may even reduce it [85, 88]. Although more research is needed to directly compare the reinforcing nature of a wider variety of HPFs to MPFs in humans, there is preliminary evidence that HPFs have a high reinforcement value (particularly for individuals prone to obesity).

In animal models, the strength of reinforcement for HPFs relative to nicotine is very clear. The majority of animal studies that administer food as a primary reinforcer use sucrose or flavored sucrose pellets, an HPF [89, 90]. From this extensive literature, we know that animals will self-administer HPF in a much wider range of conditions than nicotine; in fact, it is often the behavior trained first before attempting to train an animal to self-administer a drug [91]. Cues presented with HPFs rapidly become secondary reinforcers and places where HPFs are administered become strongly preferred [92–94]. The ability of HPFs to rapidly deliver refined carbohydrates, fat and sweet tastes appears to play a key role in their reinforcing nature, as these factors are all highly reinforcing even when studied in isolation [95–97]. Animals will self-administer sweet tastes over cocaine more than 80% of the time [98, 99]. In contrast, animals choose to self-administer nicotine over cocaine less than 20% of the time [100]. While it is clear that animals will overwhelmingly self-administer HPFs over standard laboratory chow (which provides complete nutrition through refined ingredients) [46, 101, 102], there is a need for animal models to directly compare the reinforcing nature of HPFs to MPFs (especially those naturally high in sugar or fat—such as fruits and nuts). However, existing research highlights the high reinforcing nature of HPFs even when compared to other addictive substances.

### Criterion 4: strong urges or cravings

In the 1988 SG report, craving for the substance was listed as a secondary criterion that was not considered necessary for identifying tobacco as addictive. However, the report states that the primary

criterion of compulsive use 'is driven by strong, often irresistible urges' [19]. In the 30 years since the report, the ability of tobacco products to trigger strong, recurrent urges or cravings has been identified as a central factor underlying their addictive nature [20]. The cues paired with tobacco use and nicotine delivery quickly become salient incentive drivers of behavior [103]. Consistent with this finding, cravings in response to tobacco-associated cues are a major driver of use in humans and urges to smoke are predictive of relapse during quit attempts [20]. Further, 'craving, or a strong desire or urge to use tobacco' is now a diagnostic indicator of a tobacco use disorder [17]. Based on this updated science, we propose that the ability of a substance to trigger strong cravings or urges should be a primary criterion for evaluating the addictiveness of a substance.

While experiencing a strong urge or craving for high-calorie food may be adaptive when calorically depleted, craving for HPFs commonly occur even when individuals are satiated [104, 105]. The most commonly craved foods are all common HPFs (e.g. chocolate, sweets, pizza) [104, 105] and the neural substrates underpinning cravings for HPFs and other addictive substances largely overlap [106]. As with tobacco, stimuli paired with HPFs become salient motivational cues and cue-induced craving for HPFs is implicated in more frequent HPF intake, loss of control over HPF intake (e.g. binge episodes), difficulty losing weight and a failure to reduce HPF intake in the face of serious health conditions [33–37, 104, 107–109]. In contrast to HPFs, cravings for MPFs (e.g. fruits and vegetables) are less frequent and are associated with positive health outcomes (e.g. higher diet quality, longer sleep duration, successful weight loss) [105, 110–112]. Thus, there is evidence that HPFs (but not MPFs) meet the criterion of triggering strong urges or cravings in a manner consistent with an addictive substance. Future research is needed to investigate whether the intensity of cravings for HPFs is comparable to that of other addictive substances, such as tobacco.

## SECONDARY CRITERIA

The SG report listed secondary criteria in their evaluation of the addictiveness of tobacco products, including withdrawal and tolerance [19]. These secondary criteria were not considered necessary, or sufficient, for determining the addictive nature of tobacco [19]. Science on tobacco addiction since the report further supports the secondary nature of withdrawal and tolerance. Many aspects of the withdrawal syndrome (e.g. difficulty concentrating and increased appetite) are only weakly related to relapse and the development of tolerance to the pleasurable effects of nicotine is limited [20]. While there is evidence of both withdrawal and tolerance in HPF intake [113–116], in parallel to the SG report, we do not include them as primary criteria.

## A ROADMAP FOR FUTURE DIRECTIONS

Based on the criteria used to establish the addictive nature of tobacco, we conclude that there is sufficient evidence that HPFs can

be considered addictive substances. It has been the *status quo* to treat industrial HPFs as food, not as the highly refined substances that they are, whose properties and components must be studied. Progress has been made identifying ingredients that amplify the effects of nicotine, such as menthol and sweeteners [117, 118]. To identify the key addictive components in HPFs, the same care and control employed in understanding the addictive potential of drugs needs to be applied to studies of HPFs.

In the case of industrial tobacco products, their complexity and inclusion of thousands of chemicals made identifying a single addictive agent challenging. In the SG report, nicotine was identified as the key addictive component in tobacco products as it was psychoactive, reinforcing and consumed compulsively [19], but at the time of the SG report (and still currently), the specific dose or absorption rate at which nicotine becomes addictive is unknown [119, 120]. The presence of nicotine alone in a product is not sufficient for it to be considered addictive. For example, some foods naturally contain low levels of nicotine but do not trigger addictive consumption (e.g. eggplant, cauliflower). Products such as nicotine patches, that deliver nicotine at lower doses and at a lower rate, also have low addictive potential. It is the ability of tobacco products such as cigarettes to deliver nicotine rapidly and in high doses that is key to their addictive potential.

Like industrial tobacco products, HPFs are complex substances that are psychoactive, highly reinforcing, strongly craved and consumed compulsively. The foods that people report being most likely to consume in an addictive manner are all HPFs that deliver both refined carbohydrates and added fats (i.e. chocolate, ice cream, French fries and pizza), followed by HPFs that contain refined carbohydrates without high levels of fat (i.e. breakfast cereal, gummy candy and soda) [14, 16]. Further, these HPFs are designed to rapidly deliver these unnaturally high doses of refined carbohydrates and fats due to significant changes to the food matrix during processing that removes ingredients that would slow down eating rate and absorption (e.g. water and fiber) [25]. These HPFs are energy-dense substances that quickly deliver bioavailable calories into the body, which then activates the reward systems through the gut-brain axis [121]. The exact dose and absorption rate at which refined carbohydrates and added fats become capable of triggering an addictive process is unknown, just as it is for nicotine [119, 120]. This is an important area of future research that may aid in the reformulation of HPFs to reduce addictive potential.

Investigating the role of additives that enhance the somatosensory properties of HPFs is also an important area of future research. In the 1970s additives to industrial cigarettes became more common, with hundreds of additives acknowledged by the tobacco industry [122]. The typical industrial cigarette in the United States is 10% additives by weight with sugar and other taste-based ingredients (e.g. menthol, cocoa, licorice) as main additives [122]. These additives enhance the somatosensory appeal, taste and smell of industrial tobacco products. Smokers develop strong brand loyalty and are reluctant to switch to other industrial brands [123] or to new products, even if they deliver high doses of nicotine (e.g. electronic cigarettes [20]). Denicotinized cigarettes that deliver the somatosensory experience of smoking, but

no nicotine, reduce cigarette craving and are even chosen over intravenous nicotine administration [124]. Further evidence for the importance of somatosensory properties is demonstrated by the recent proposal to remove menthol-flavored cigarettes and flavored cigars from the market-place in the United States. The change to the somatosensory profile is estimated to lead hundreds of thousands of people to quit smoking or to never initiate smoking, despite no reduction in the nicotine levels in the products [125]. Thus, tobacco products are addictive, not just because of the existence of nicotine, but because they are industrial products that have been optimized to deliver nicotine rapidly at high doses and are enhanced through additives to have unique somatosensory signatures.

Industrial HPFs are also optimized through additives that pair specific somatosensory properties (e.g. taste, smell, texture and mouth-feel) with the delivery of high doses of refined carbohydrates and added fats [25]. These additives are unlikely to be reinforcing on their own, but when combined with refined carbohydrates and added fats they probably play a key role in amplifying the addictive nature of industrial HPFs. Of concern, many industrial additives (e.g. flavor enhancers such as nucleotides and artificial flavors, texturizers such as emulsifiers) are not typically used in home cooking. Many somatosensory properties that become powerful drivers of intake (e.g. artificial cheese flavoring, chewiness of gummy candy) are typically only available in industrial HPFs. As with personal tobacco product preferences [123], individual differences in the experience of specific somatosensory profiles paired with the delivery of rapidly absorbed carbohydrates and/or added fats may contribute to personal preferences or brand loyalty for specific HPFs. There are also additives, such as non-nutritive sweeteners, that mimic the effects of sugar by intensely activating sweet taste receptors on the tongue without delivering calories. Non-nutritive sweeteners are reinforcing and increase activity in reward centers of the brain (although possibly to a lesser degree than nutritive sweeteners) [126]. Non-nutritive sweeteners are frequently included in HPFs alongside refined carbohydrates and/or fat [127], but some products (such as diet sodas) include only non-nutritive sweeteners that are often combined with reinforcing ingredients such as caffeine. While some randomized control trials have found limited effects of non-nutritive sweeteners on energy intake or body weight, overall there are numerous gaps in the evidence based on the impact of non-nutritive sweeteners on health outcomes [128]. Thus, the understanding of the addictive potential of non-nutritive sweeteners (when consumed alongside caloric ingredients and when amplified by additives such as caffeine) is an important area of future study.

Another important question is whether certain MPFs (or MPF combinations) can trigger addictive responses in some circumstances. Although some MPFs have naturally high levels of carbohydrates (e.g. fruit) or fat (e.g. salmon), people report that these foods are unlikely to be consumed in an addictive manner [1, 14–16, 42]. As reviewed above, MPFs (such as fruits and vegetables) are rarely consumed compulsively, are less prone to reinforcement sensitization and cravings for MPFs are associated with better health [42–45, 80, 84, 85, 105, 110–112]. However, MPFs are also diverse. Naturally high in fat MPFs that typically also have high levels of added sodium (e.g.

bacon, cheese, salted nuts) are associated with higher subjective ratings of loss-of-control, craving and enjoyment relative to other MPFs (e.g. avocados, eggs, apples, bananas) [16]. Thus, it is important to investigate whether high-fat MPFs with high levels of added sodium have addictive properties. Additionally, MPFs may be altered and combined in ways that may increase their addictive potential. For example, the blending together of high-sugar fruits and high-fat yogurt into a smoothie creates an MPF that delivers high levels of rapidly absorbed carbohydrates combined with fat akin to a HPF. It will be important for future research to investigate the boundary at which some MPFs (or MPF combinations) may meet the criteria for addictive substances based on the proposed criteria.

## CONCLUSION

Here, we provide evidence that HPFs meet the three criteria of an addictive substance as outlined by the 1988 SG report on the addictive nature of tobacco products. Like tobacco, HPFs (1) trigger compulsive use, (2) have psychoactive effects and (3) are highly reinforcing [19]. In addition, HPFs (4) trigger strong urges and cravings, which has emerged as a strong predictor of addictive use [20]. These HPFs are highly complex substances that cannot be simplified to a single chemical agent acting through a specific central mechanism. This is also true of industrial tobacco products, which contain thousands of chemicals (in addition to nicotine), an optimized pharmacokinetic profile, and unique somatosensory properties that enhance their addictive nature [122]. A dose and rate profile of a single addictive chemical was not used to identify tobacco products as addictive. However, HPFs meet the actual scientific criteria used to determine that tobacco products are addictive. More research into the exact properties and components is needed, but their ability to rapidly deliver high doses of refined carbohydrates and/or fat appear key to their addictive potential. Thus, we conclude that HPFs can be considered addictive substances based on scientifically established criteria.

The scientific evidence that tobacco products are addictive supported the advancement of new treatment, policy and public health approaches that helped significantly reduce their use. We must consider the huge public health costs of misclassifying an addictive substance as non-addictive. When tobacco was misclassified this led the public to be ill-informed about the risks of tobacco products, reduced the likelihood that addiction mechanisms were targeted in treatments and allowed the industry to develop more addictive products with limited oversight and to target vulnerable populations, including children and racial and ethnic minorities [29]. Identifying tobacco products not just as unhealthy but also as addictive undermined the industry's claim that smoking was solely an act of free will and an 'adult choice' [129]. The SG report supported increased scrutiny of industry practices, contributed to the success of legal proceedings against the tobacco industry and increased public support for policy initiatives that aimed to protect children (e.g. restrictions in tobacco marketing to minors) and alter the tobacco environment (e.g. taxation and reducing vending machine access) [29].

Tobacco companies (i.e. Philip Morris and R.J. Reynolds) bought food and beverage companies (e.g. Kraft General Foods, Nabisco) and became some of the biggest producers of HPFs in the world from 1980 to the mid-2000s [130, 131]. Tobacco industry documents demonstrate that they applied their playbook from tobacco to increase the profits from their food and beverage portfolios, including the use of flavored additives and marketing strategies to target children and racial/ethnic minorities [130, 131]. Currently, poor diets dominated by HPFs are contributing to preventable deaths to a comparable degree as tobacco products [132]. Understanding whether addiction contributes to HPF intake may lead to new treatments, as preliminary research finds that behavioral and pharmacological interventions that target addictive mechanisms may reduce compulsive HPF intake [133, 134]. Importantly, if the science supports that HPFs are not just unhealthy, but addictive, this challenges the assertion that excessive HPF intake is purely a matter of choice. This may increase scrutiny of industrial practices in the development and marketing of HPFs (particularly to children). At the time of the SG report, tobacco products were the largest cause of preventable death; however, there was still some resistance to accepting their addictive and harmful nature. This was due in large part to industry efforts to undermine confidence in the science on the addictive nature of tobacco [29], and because tobacco products were 'so familiar they [had] lost their saliency' [135]. This delayed the implementation of effective strategies to address this public health crisis, which cost millions of lives. Unlike smoking, we all need to eat. In the past 40 years, HPFs have become familiar substances that dominate the food environment, but we cannot lose the saliency of their potential to be addictive and harmful.

## ACKNOWLEDGEMENTS

None.

## DECLARATION OF INTERESTS

None.

## AUTHOR CONTRIBUTIONS

**Ashley N. Gearhardt:** Conceptualization (equal); writing – original draft (lead); writing – review and editing (equal). **Alexandra G. DiFelicantonio:** Conceptualization (equal); writing – original draft (supporting); writing – review and editing (equal).

## ORCID

Ashley N. Gearhardt  <https://orcid.org/0000-0003-3843-5731>

## REFERENCES

- Gearhardt AN, Schulte EM. Is food addictive? A review of the science. *Annu Rev Nutr.* 2021;41:387–410.
- Praxedes DR, Silva-Júnior AE, Macena ML, Oliveira AD, Cardoso KS, Nunes LO, et al. Prevalence of food addiction determined by the Yale Food Addiction Scale and associated factors: a systematic review with meta-analysis. *Eur Eat Disord Rev.* 2022;30:85–95.
- Yekaninejad MS, Badrooj N, Vosoughi F, Lin CY, Potenza MN, Pakpour AH. Prevalence of food addiction in children and adolescents: a systematic review and meta-analysis. *Obes Rev.* 2021; 22:e13183.
- Fletcher PC, Kenny PJ. Food addiction: a valid concept? *Neuropsychopharmacology.* 2018;43(2513):2506.
- Hebebrand J, Gearhardt AN. The concept of 'food addiction' helps inform the understanding of overeating and obesity: NO. *Am J Clin Nutr.* 2021;113:268–73.
- O'Connor RM, Kenny PJ. Utility of 'substance use disorder' as a heuristic for understanding overeating and obesity. *Progr Neuro-Psychopharmacol Biol Psychiatry.* 2022;118:110580.
- Finlayson G. Food addiction and obesity: unnecessary medicalization of hedonic overeating. *Nat Rev Endocrinol.* 2017;13:493–8.
- Fielding-Singh P, Patel ML, King AC, Gardner CD. Baseline psychosocial and demographic factors associated with study attrition and 12-month weight gain in the DIETFITS trial. *Obesity.* 2019;27:1997–2004.
- Minhas M, Murphy CM, Balodis IM, Samokhvalov AV, MacKillop J. Food addiction in a large community sample of Canadian adults: prevalence and relationship with obesity, body composition, quality of life and impulsivity. *Addiction.* 2021;116:2870–9.
- Hebebrand J, Albayrak Ö, Adan R, Antel J, Dieguez C, de Jong J, et al. 'Eating addiction', rather than 'food addiction', better captures addictive-like eating behavior. *Neurosci Biobehav Rev.* 2014;47: 295–306.
- Breslin PA. An evolutionary perspective on food and human taste. *Curr Biol.* 2013;23:R409–18.
- Musser JM, Schippers KJ, Nickel M, Mizzon G, Kohn AB, Pape C, et al. Profiling cellular diversity in sponges informs animal cell type and nervous system evolution. *Science.* 2021;374:717–23.
- Hernandez L, Hoebel BG. Food reward and cocaine increase extracellular dopamine in the nucleus accumbens as measured by microdialysis. *Life Sci.* 1988;42:1705–12.
- Schulte EM, Avena NM, Gearhardt AN. Which foods may be addictive? The roles of processing, fat content, and glycemic load. *PLOS One.* 2015;10:e0117959.
- Gearhardt AN, Hebebrand J. The concept of 'food addiction' helps inform the understanding of overeating and obesity: YES. *Am J Clin Nutr.* 2021;113:263–7.
- Schulte EM, Smeal JK, Gearhardt AN. Foods are differentially associated with subjective effect report questions of abuse liability. *PLOS One.* 2017;12:e0184220.
- American Psychiatric Association (APA). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. Washington, DC: APA; 2013.
- Gearhardt AN, Hebebrand J. The concept of 'food addiction' helps inform the understanding of overeating and obesity: debate consensus. *Am J Clin Nutr.* 2021;113:274–6.
- US Department of Health and Human Services (USDHHS). *Health Consequences of Smoking: Nicotine Addiction. A report of the Surgeon General 1988*. Washington, DC: USDHHS, Office of Smoking and Health; 1988.
- West R, Cox S. The 1988 US Surgeon General's report *Nicotine Addiction: how well has it stood up to three more decades of research?* *Addiction.* 2022;117(8):2346–2350. <https://doi.org/10.1111/add.15754>
- Poti JM, Mendez MA, Ng SW, Popkin BM. Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *Am J Clin Nutr.* 2015;101: 1251–62.
- Fazzino TL. The reinforcing natures of hyper-palatable foods: behavioral evidence for their reinforcing properties and the role of the US food industry in promoting their availability. *Curr Addict Rep.* 2022; 1–9. <https://doi.org/10.1007/s40429-022-00417-8>
- Monteiro CA, Astrup A. Does the concept of 'ultra-processed foods' help inform dietary guidelines, beyond conventional classification systems? YES. *Am J Clin Nutr.* 2022;nqac122. <https://doi.org/10.1093/ajcn/nqac122>

24. Astrup A, Monteiro C. Does the concept of 'ultra-processed foods' help inform dietary guidelines, beyond conventional classification systems? *NO. Am J Clin Nutr.* 2022;nqac123. <https://doi.org/10.1093/ajcn/nqac123>
25. Monteiro CA, Cannon G, Levy RB, Moubarac J-C, Louzada ML, Rauber F, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr.* 2019;22:936–41.
26. Monteiro CA, Cannon GJ. The role of the transnational ultra-processed food industry in the pandemic of obesity and its associated diseases: problems and solutions. *World Nutrition.* 2019;10:89–99.
27. Cotter T, Kotov A, Wang S, Murukutla N. 'Warning: Ultra-processed'—a call for warnings on foods that aren't really foods. *BMJ Glob Health.* 2021;6:e007240.
28. Cross GS, Proctor RN, Proctor R. *Packaged Pleasures: How Technology and Marketing Revolutionized Desire.* Chicago, IL: University of Chicago Press; 2014.
29. Brownell KD, Warner KE. The perils of ignoring history: big tobacco played dirty and millions died. How similar is big food? *Milbank Q.* 2009;87:259–94.
30. Burns DM, Lee L, Shen LZ, Gilpin E, Tolley HD, Vaughn J et al. Cigarette smoking behavior in the United States. Changes in cigarette-related disease risks and their implication for prevention and control. *Smoking and Tobacco Control Monograph no. 8.* Bethesda, MD: US Department of Health and Human Services; 1997, pp. 13–42.
31. Casagrande SS, Wang Y, Anderson C, Gary TL. Have Americans increased their fruit and vegetable intake? The trends between 1988 and 2002. *Am J Prev Med.* 2007;32:257–63.
32. Anthony JC, Warner LA, Kessler RC. Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants: basic findings from the National Comorbidity Survey. *Exp Clin Psychopharmacol.* 1994;2:244.
33. Ponso V, Rosato R, Tarsia E, Goitre I, De Michieli F, Fadda M, et al. Self-reported adherence to diet and preferences towards type of meal plan in patient with type 2 diabetes mellitus. A cross-sectional study. *Nutr Metab Cardiovasc Dis.* 2017;27:642–50.
34. Shultz JA, Sprague MA, Branan LJ, Lambeth S. A comparison of views of individuals with type 2 diabetes mellitus and diabetes educators about barriers to diet and exercise. *J Health Commun.* 2001;6:99–115.
35. White S, Bissell P, Anderson C. A qualitative study of cardiac rehabilitation patients' perspectives on making dietary changes. *J Hum Nutr Diet.* 2011;24:122–7.
36. Ebrahim Z, Ahmed T, De Villiers A. Factors influencing adherence to dietary guidelines: a qualitative study on the experiences of patients with type 2 diabetes attending a clinic in Cape Town. *J Endocrinol Metab Diab South Africa.* 2014;19:76–84.
37. LaCharity LA, King KM, King KB. The experiences of postmenopausal women with coronary artery disease. *West J Nurs Res.* 1997;19:583–607.
38. Budak AR, Thomas SE. Food craving as a predictor of 'relapse' in the bariatric surgery population: a review with suggestions. *Bariatric Nurs Surg Patient Care.* 2009;4:115–21.
39. Smith KR, Aghababian A, Papanoni A, Veldhuizen MG, Kamath V, Harris C, et al. One year follow-up of taste-related reward associations with weight loss suggests a critical time to mitigate weight regain following bariatric surgery. *Nutrients.* 2021;13:3943.
40. Li Z, Pan Y, Zhang Y, Qin J, Lei X. Dietary experiences after bariatric surgery in patients with obesity: a qualitative systematic review. *Obes Surg.* 2022;32:2023–34.
41. Davis C. Compulsive overeating as an addictive behavior: overlap between food addiction and binge eating disorder. *Curr Obes Rep.* 2013;2:171–17.
42. Ayton A, Ibrahim A, Dugan J, Galvin E, Wright OW. Ultra-processed foods and binge eating: a retrospective observational study. *Nutrition.* 2021;84:111023.
43. Rosen JC, Leitenberg H, Fisher C, Khazam C. Binge-eating episodes in bulimia nervosa: the amount and type of food consumed. *Int J Eat Disord.* 1986;5:255–67.
44. Allison S, Timmerman GM. Anatomy of a binge: food environment and characteristics of nonpurge binge episodes. *Eat Behav.* 2007;8:31–8.
45. Bertoli S, Spadafranca A, Bes-Rastrollo M, Martinez-Gonzalez MA, Ponissi V, Beggio V, et al. Adherence to the Mediterranean diet is inversely related to binge eating disorder in patients seeking a weight loss program. *Clin Nutr.* 2015;34:107–14.
46. Johnson PM, Kenny PJ. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nat Neurosci.* 2010;13:635–41.
47. Krasnova IN, Marchant NJ, Ladenheim B, McCoy MT, Panlilio LV, Bossert JM, et al. Incubation of methamphetamine and palatable food craving after punishment-induced abstinence. *Neuropsychopharmacology.* 2014;39:2008–16.
48. Soria R, Stapleton JM, Gilson SF, Sampson-Cone A, Henningfield JE, London ED. Subjective and cardiovascular effects of intravenous nicotine in smokers and non-smokers. *Psychopharmacology.* 1996;128:221–6.
49. Wills L, Kenny PJ. Addiction-related neuroadaptations following chronic nicotine exposure. *J Neurochem.* 2021;157:1652–73.
50. Heinz A, Daedelow LS, Wackerhagen C, Di Chiara G. Addiction theory matters—why there is no dependence on caffeine or antidepressant medication. *Addict Biol.* 2020;25:e12735.
51. Di Chiara G, Imperato A. Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci USA.* 1988;85:5274–8.
52. Bassareo V, De Luca MA, Aresu M, Aste A, Ariu T, Di Chiara G. Differential adaptive properties of accumbens shell dopamine responses to ethanol as a drug and as a motivational stimulus. *Eur J Neurosci.* 2003;17:1465–72.
53. Nasser JA, Bradley LE, Leitzsch JB, Chohan O, Fasulo K, Haller J, et al. Psychoactive effects of tasting chocolate and desire for more chocolate. *Physiol Behav.* 2011;104:117–21.
54. Casperson SL, Lanza L, Albajri E, Nasser JA. Increasing chocolate's sugar content enhances its psychoactive effects and intake. *Nutrients.* 2019;11:596.
55. Spring B, Schneider K, Smith M, Kendzor D, Appelhans B, Hedeker D, et al. Abuse potential of carbohydrates for overweight carbohydrate cravers. *Psychopharmacology.* 2008;197:637–47.
56. Cummings JR, Schiestl ET, Tomiyama AJ, Mamtara T, Gearhardt AN. Highly processed food intake and immediate and future emotions in everyday life. *Appetite.* 2022;169:105868.
57. Schulte EM, Sonnevill KR, Gearhardt AN. Subjective experiences of highly processed food consumption in individuals with food addiction. *Psychol Addict Behav.* 2019;33:144.
58. Joyner MA, Schulte EM, Wilt AR, Gearhardt AN. Addictive-like eating mediates the association between eating motivations and elevated body mass index. *Transl Issues Psychol Sci.* 2015;1:217.
59. Hajnal A, Smith GP, Norgren R. Oral sucrose stimulation increases accumbens dopamine in the rat. *Am J Physiol Reg Integr Comparat Physiol.* 2004;286:R31–7.
60. Bassareo V, Di Chiara G. Differential influence of associative and nonassociative learning mechanisms on the responsiveness of prefrontal and accumbal dopamine transmission to food stimuli in rats fed *ad libitum*. *J Neurosci.* 1997;17:851–61.
61. Bassareo V, De Luca MA, Di Chiara G. Differential expression of motivational stimulus properties by dopamine in nucleus accumbens shell versus core and prefrontal cortex. *J Neurosci.* 2002;22:4709–19.
62. De Luca MA. Habituation of the responsiveness of mesolimbic and mesocortical dopamine transmission to taste stimuli. *Front Integr Neurosci.* 2014;8:21.

63. Tellez LA, Ren X, Han W, Medina S, Ferreira JG, Yeckel CW, et al. Glucose utilization rates regulate intake levels of artificial sweeteners. *J Physiol*. 2013;591:5727–44.
64. Tellez LA, Medina S, Han W, Ferreira JG, Licona-Limón P, Ren X, et al. A gut lipid messenger links excess dietary fat to dopamine deficiency. *Science*. 2013;341:800–2.
65. Alhadeff AL, Goldstein N, Park O, Klima ML, Vargas A, Betley JN. Natural and drug rewards engage distinct pathways that converge on coordinated hypothalamic and reward circuits. *Neuron*. 2019;103:891–908, e6.
66. Dayan P. 'Liking' as an early and editable draft of long-run affective value. *PLOS Biol*. 2022;20:e3001476.
67. DiFeliceantonio AG, Coppin G, Rigoux L, Thanarajah SE, Dagher A, Tittgemeyer M, et al. Supra-additive effects of combining fat and carbohydrate on food reward. *Cell Metab*. 2018;28:33–44.
68. Caref K, Nicola SM. Endogenous opioids in the nucleus accumbens promote approach to high-fat food in the absence of caloric need. *eLife*. 2018;7:e34955.
69. Alsiö J, Olszewski PK, Levine AS, Schiöth HB. Feed-forward mechanisms: addiction-like behavioral and molecular adaptations in overeating. *Front Neuroendocrinol*. 2012;33:127–39.
70. Drewnowski A, Krahn DD, Demitrack MA, Nairn K, Gosnell BA. Naloxone, an opiate blocker, reduces the consumption of sweet high-fat foods in obese and lean female binge eaters. *Am J Clin Nutr*. 1995;61:1206–12.
71. Yeomans MR, Gray RW. Opioid peptides and the control of human ingestive behaviour. *Neurosci Biobehav Rev*. 2002;26:713–28.
72. Giraud S, Grace M, Welch C, Billington CJ, Levine AS. Naloxone's anorectic effect is dependent upon the relative palatability of food. *Pharmacol Biochem Behav*. 1993;46:917–21.
73. Joyner MA, Kim S, Gearhardt AN. Investigating an incentive-sensitization model of eating behavior: impact of a simulated fast-food laboratory. *Clin Psychol Sci*. 2017;5:1014–26.
74. Meule A, Kübler A. Food cravings in food addiction: the distinct role of positive reinforcement. *Eat Behav*. 2012;13:252–5.
75. Pomerleau OF. Individual differences in sensitivity to nicotine: implications for genetic research on nicotine dependence. *Behav Genet*. 1995;25:161–77.
76. Feig EH, Piers AD, Kral TV, Lowe MR. Eating in the absence of hunger is related to loss-of-control eating, hedonic hunger, and short-term weight gain in normal-weight women. *Appetite*. 2018;123:317–24.
77. Lansigan RK, Emond JA, Gilbert-Diamond D. Understanding eating in the absence of hunger among young children: a systematic review of existing studies. *Appetite*. 2015;85:36–47.
78. Kral TV, Moore RH, Chittams J, O'Malley L, Jones E, Quinn RJ, et al. Does eating in the absence of hunger extend to healthy snacks in children? *Pediatr Obes*. 2020;15:e12659.
79. Epstein LH, Salvy SJ, Carr KA, Dearing KK, Bickel WK. Food reinforcement, delay discounting and obesity. *Physiol Behav*. 2010;100:438–45.
80. McCullough MB, Guilkey H, Stark L. Cookie or fruit? Relative reinforcing value of snack foods among preschoolers with overweight/obesity compared to healthy weight. *Appetite*. 2017;111:187–94.
81. Epstein LH, Leddy JL, Temple JL, Faith MS. Food reinforcement and eating: a multilevel analysis. *Psychol Bull*. 2007;133:884.
82. Temple JL, Bulkley AM, Badawy RL, Krause N, McCann S, Epstein LH. Differential effects of daily snack food intake on the reinforcing value of food in obese and nonobese women. *Am J Clin Nutr*. 2009;90:304–13.
83. Temple JL, Legierski CM, Giacomelli AM, Salvy S-J, Epstein LH. Overweight children find food more reinforcing and consume more energy than do nonoverweight children. *Am J Clin Nutr*. 2008;87:1121–7.
84. Clark EN, Dewey AM, Temple JL. Effects of daily snack food intake on food reinforcement depend on body mass index and energy density. *Am J Clin Nutr*. 2010;91:300–8.
85. Temple JL, Ziegler AM, Crandall AK, Mansouri T, Hatzinger L, Barich R, et al. Sensitization of the reinforcing value of high energy density foods is associated with increased zBMI gain in adolescents. *Int J Obes*. 2022;46:581–7.
86. Temple JL, Ziegler AM, Crandall AK, Mansouri T, Epstein LH. Sensitization of the reinforcing value of food: a novel risk factor for overweight in adolescents. *Int J Obes*. 2020;44:1918–27.
87. Temple JL. Behavioral sensitization of the reinforcing value of food: What food and drugs have in common. *Prev Med*. 2016;92:90–9.
88. Casperson SL, Jahns L, Temple JL, Appleton KM, Duke SE, Roemmich JN. Consumption of a variety of vegetables to meet dietary guidelines for Americans' recommendations does not induce sensitization of vegetable reinforcement among adults with overweight and obesity: a randomized controlled trial. *J Nutr*. 2021;151:1665–72.
89. Fisher Scientific. Bio-Serv™ Dustless Precision Pellets™ Grain-Based Rodent Diet. Available from: <https://www.fishersci.com/shop/products/bio-serv-dustless-precision-pellets-grain-based-rodent-diet-7/p-4185725>. Accessed 9 March 2022
90. TestDiet. TestDiet Tablet Spec Sheet. Available from: [https://www.testdiet.com/cs/groups/lolweb/@testdiet/documents/web\\_content/mdrf/mdi2/~edisp/ducm04\\_026412.pdf](https://www.testdiet.com/cs/groups/lolweb/@testdiet/documents/web_content/mdrf/mdi2/~edisp/ducm04_026412.pdf). Accessed 9 March 2022
91. Elayouby KS, Ishikawa M, Dukes AJ, Smith AC, Lu Q, Fowler CD, et al.  $\alpha 3^*$  Nicotinic acetylcholine receptors in the habenula-interpeduncular nucleus circuit regulate nicotine intake. *J Neurosci*. 2021;41:1779–87.
92. Peciña S, Berridge KC. Dopamine or opioid stimulation of nucleus accumbens similarly amplify cue-triggered 'wanting' for reward: entire core and medial shell mapped as substrates for PIT enhancement. *J Neurosci*. 2013;37:1529–40.
93. King CP, Tripi JA, Hughson AR, Horvath AP, Lamparelli AC, Holl KL, et al. Sensitivity to food and cocaine cues are independent traits in a large sample of heterogeneous stock rats. *Sci Rep*. 2021;11:1–17.
94. Berridge KC, Ho C-Y, Richard JM, DiFeliceantonio AG. The tempted brain eats: pleasure and desire circuits in obesity and eating disorders. *Brain Res*. 2010;1350:43–64.
95. Lucas F, Sclafani A. Differential reinforcing and satiating effects of intragastric fat and carbohydrate infusions in rats. *Physiol Behav*. 1999;66:381–8.
96. Elizalde G, Sclafani A. Fat appetite in rats: flavor preferences conditioned by nutritive and non-nutritive oil emulsions. *Appetite*. 1990;15(3):189–97.
97. Ackroff K, Dym C, Yiin Y-M, Sclafani A. Rapid acquisition of conditioned flavor preferences in rats. *Physiol Behav*. 2009;97:406.
98. Lenoir M, Serre F, Cantin L, Ahmed SH. Intense sweetness surpasses cocaine reward. *PLOS One*. 2007;2:e698.
99. Ahmed SH, Guillem K, Vandaele Y. Sugar addiction: pushing the drug-sugar analogy to the limit. *Curr Opin Clin Nutr Metab Care*. 2013;16:434–9.
100. Manzardo A, Stein L, Belluzzi J. Rats prefer cocaine over nicotine in a two-lever self-administration choice test. *Brain Res*. 2002;924:10–9.
101. DiFeliceantonio AG, Mabrouk OS, Kennedy RT, Berridge KC. Enkephalin surges in dorsal neostriatum as a signal to eat. *Curr Biol*. 2012;22:1918–24.
102. DiFeliceantonio AG, Berridge KC. Dorsolateral neostriatum contribution to incentive salience: opioid or dopamine stimulation makes one reward cue more motivationally attractive than another. *Eur J Neurosci*. 2016;43:1203–18.
103. Caggiula AR, Donny EC, White AR, Chaudhri N, Booth S, Gharib MA, et al. Cue dependency of nicotine self-administration and smoking. *Pharmacol Biochem Behav*. 2001;70:515–30.

104. Meule A. Twenty years of the food cravings questionnaires: a comprehensive review. *Curr Addict Rep.* 2020;7:30–43.
105. Meule A. The psychology of food cravings: the role of food deprivation. *Curr Nutr Rep.* 2020;9:251–7.
106. Noori HR, Linan AC, Spanagel R. Largely overlapping neuronal substrates of reactivity to drug, gambling, food and sexual cues: a comprehensive meta-analysis. *Eur Neuropsychopharmacol.* 2016;26:1419–30.
107. Veilleux JC, Skinner KD. Smoking, food, and alcohol cues on subsequent behavior: a qualitative systematic review. *Clin Psychol Rev.* 2015;36:13–27.
108. Boswell RG, Kober H. Food cue reactivity and craving predict eating and weight gain: a meta-analytic review. *Obes Rev.* 2016;17:159–77.
109. Greeno CG, Wing RR, Shiffman S. Binge antecedents in obese women with and without binge eating disorder. *J Consult Clin Psychol.* 2000;68:95.
110. Richard A, Meule A, Reichenberger J, Blechert J. Food cravings in everyday life: an EMA study on snack-related thoughts, cravings, and consumption. *Appetite.* 2017;113:215–23.
111. Anton SD, Gallagher J, Carey VJ, Laranjo N, Cheng J, Champagne CM, et al. Diet type and changes in food cravings following weight loss: findings from the Pounds Lost Trial. *Eat Weight Disord Stud Anorexia Bulimia Obes.* 2012;17:e101–8.
112. Kracht CL, Chaput J-P, Martin CK, Champagne CM, Katzmarzyk PT, Staiano AE. Associations of sleep with food cravings, diet, and obesity in adolescence. *Nutrients.* 2019;11:2899.
113. Schulte EM, Smeal JK, Lewis J, Gearhardt AN. Development of the highly processed food withdrawal scale. *Appetite.* 2018;131:148–54.
114. Avena NM, Rada P, Hoebel BG. Evidence for sugar addiction: behavioral and neurochemical effects of intermittent, excessive sugar intake. *Neurosci Biobehav Rev.* 2008;32:20–39.
115. Papantoni A, Shearrer G, Sadler J, Burger K (Eds). *Obesity: High Sugar Intake From Beverages is Associated With Decreased Dopaminergic Brain Response to Sugar.* Hoboken, NJ: Wiley; 2021.
116. Parnarouskis L, Gearhardt AN. Preliminary evidence that tolerance and withdrawal occur in response to ultra-processed foods. *Curr Addict Rep.* 2022;1–8. <https://doi.org/10.1007/s40429-022-00425-8>
117. Zhang M, Harrison E, Biswas L, Tran T, Liu X. Menthol facilitates dopamine-releasing effect of nicotine in rat nucleus accumbens. *Pharmacol Biochem Behav.* 2018;175:47–52.
118. Kroemer NB, Veldhuizen MG, Delvy R, Patel BP, O'Malley SS, Small DM. Sweet taste potentiates the reinforcing effects of e-cigarettes. *Eur Neuropsychopharmacol.* 2018;28:1089–102.
119. Benowitz NL, Hukkanen J, Jacob P III. Nicotine Chemistry, Metabolism, Kinetics and Biomarkers. *Handb Exp Pharmacol.* 2009;19:29–60.
120. Benowitz NL, Henningfield JE. Establishing a nicotine threshold for addiction—the implications for tobacco regulation. *N Engl J Med.* 1994;14(331):123–5.
121. Kelly A, Baugh ME, Oster ME, DiFeliceantonio AG. The impact of caloric availability on eating behavior and ultra-processed food reward. *Appetite.* 2022;78:106274.
122. Rabinoff M, Caskey N, Rissling A, Park C. Pharmacological and chemical effects of cigarette additives. *Am J Public Health.* 2007;97:1981–91.
123. DeCicca P, Kenkel D, Liu F, Somerville J. Quantifying brand loyalty: evidence from the cigarette market. *J Health Econ.* 2021;79:102512.
124. Rose JE, Salley A, Behm FM, Bates JE, Westman EC. Reinforcing effects of nicotine and non-nicotine components of cigarette smoke. *Psychopharmacology.* 2010;210:1–12.
125. Cadham CJ, Sanchez-Romero LM, Fleischer NL, Mistry R, Hirschtick JL, Meza R, et al. The actual and anticipated effects of a menthol cigarette ban: a scoping review. *BMC Public Health.* 2020;20:1–17.
126. Yunker AG, Patel R, Page KA. Effects of non-nutritive sweeteners on sweet taste processing and neuroendocrine regulation of eating behavior. *Curr Nutr Rep.* 2020;9:278–89.
127. Dunford EK, Taillie LS, Miles DR, Eyles H, Tolentino-Mayo L, Ng SW. Non-nutritive sweeteners in the packaged food supply—an assessment across 4 countries. *Nutrients.* 2018;10:257.
128. Lohner S, Toews I, Meerpohl JJ. Health outcomes of non-nutritive sweeteners: analysis of the research landscape. *Nutr J.* 2017;16:1–21.
129. Henningfield JE, Rose CA, Zeller M. Tobacco industry litigation position on addiction: continued dependence on past views. *Tob Control.* 2006;15:iv27–36.
130. Nguyen KH, Glantz SA, Palmer CN, Schmidt LA. Tobacco industry involvement in children's sugary drinks market. *BMJ.* 2019;364:1736.
131. Nguyen KH, Glantz SA, Palmer CN, Schmidt LA. Transferring racial/ethnic marketing strategies from tobacco to food corporations: Philip Morris and Kraft General Foods. *Am J Public Health.* 2020;110:329–36.
132. Bauer UE, Briss PA, Goodman RA, Bowman BA. Prevention of chronic disease in the 21st century: Elimination of the leading preventable causes of premature death and disability in the USA. *Lancet.* 2014;384:45–52.
133. Carbone EA, Caroleo M, Rania M, Calabrò G, Staltari FA, de Filippis R, et al. An open-label trial on the efficacy and tolerability of naltrexone/bupropion SR for treating altered eating behaviours and weight loss in binge eating disorder. *Eat Weight Disord Stud Anorexia Bulimia Obes.* 2021;26:779–88.
134. Kwag KH, Kim YR, Lee YK, Lee HR, Lee JY, Lee J, et al. Feasibility and acceptability of a digital tele-guided intervention targeting components of the addictive appetite model for bulimia nervosa and binge-eating disorder in Korea. *Int J Eating Disord.* 2022;55:977–82.
135. Pollin W. The role of the addictive process as a key step in causation of all tobacco-related diseases. *JAMA.* 1984;252:2874.

**How to cite this article:** Gearhardt AN, DiFeliceantonio AG. Highly processed foods can be considered addictive substances based on established scientific criteria. *Addiction.* 2023;118(4):589–98. <https://doi.org/10.1111/add.16065>